



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Death rate of Berlin compared with other cities.

I have the honor to report that the death rate of Berlin for the week ended September 12 was lower than it has been since the middle of July, amounting, calculated on the year, to 16 per thousand of the population, being thereby, however, considerably higher than for the corresponding week of last year, in which it amounted to 14 per thousand. Of the large German cities during this week only the following showed more favorable figures than Berlin, viz, Hamburg, Barmen, Frankfort-on-the-Main, Stuttgart, Hanover, and Charlottenburg (with 10.5). The death rate of Schöneberg (17 per thousand), Karlsruhe, Munich, Cologne, and Magdeburg was not much higher. On the other hand, the death rate of Breslau, Nuremberg, Leipzig, Dresden, and Stettin was considerably higher than the Berlin rate, the death rate of Vienna was somewhat higher than that of Berlin, while the rates of London and Paris were lower than the Berlin figure.

The decrease in the number of deaths was confined exclusively to children in the first year of life. The infant death rate fell from 7.2 per year and mille to 6 in the week under discussion, being lower than that of Munich and Hamburg and not even the half of the Breslau, Leipzig, and Nuremberg rate. Acute diseases of the respiratory organs showed again a slight increase, causing 42 deaths. Acute intestinal diseases showed a decrease, claiming 119 victims (as against 168 in the foregoing week). Phthisis pulmonalis caused 61 deaths, and 51 persons succumbed to cancer. There further occurred 2 deaths from scarlet fever, 2 deaths from diphtheria, 1 death from measles, 3 deaths from typhus, and 23 persons died by violence.

GUATEMALA.

Report from Livingston, fruit port.

Acting Assistant Surgeon R. H. Peters reports as follows for the week ended September 28, 1903: Present officially estimated population, about 3,500; prevailing diseases, malarial; general sanitary condition of this port and the surrounding country during the week, good.

Bills of health were issued to the following vessels:

Date.	Name of vessel.	Number of crew.	Number of passengers from this port.	Number of passengers in transit.	Pieces of baggage disinfected.
Sept. 22	Spero	17	0	0	0
22	Anselm	41	2	0	11

The steamship *Anselm* cleared from Puerto Barrios.

HAWAIIAN ISLANDS.

Outgoing quarantine transactions at Honolulu and Hilo.

Chief Quarantine Officer Cofer reports as follows:

Outgoing quarantine transactions at this port for the week ended September 26, 1903, were as follows:

Vessels inspected and bills of health issued	7
Vessels disinfected	4